

The Cambrian Hopf Algebra


G. Châtel

July 10th, 2015


Joint work with V. Pilaud

Combinatorics


permutations



binary trees





binary sequences



Algebra

Malvenuto-Reutenauer algebra
 $\text{FQSym} = \text{vect} \langle \mathbb{F}_\tau \mid \tau \in \mathfrak{S} \rangle$ Loday-Ronco algebra
 $\text{PBT} = \text{vect} \langle \mathbb{P}_T \mid T \in \mathcal{BT} \rangle$ Solomon algebra
 $\text{Rec} = \text{vect} \langle \mathbb{X}_\eta \mid \eta \in \pm^* \rangle$

Geometry



1 Combinatorics

- Binary trees
- Cambrian trees
- Cambrian lattices

2 Algebra

- FQSym
- The Cambrian algebra

1 Combinatorics


- Binary trees
- Cambrian trees
- Cambrian lattices

2 Algebra




- FQSym
- The Cambrian algebra

Binary trees

Binary search tree = directed and labeled tree such that




increasing tree = directed and labeled tree such that labels increase along arcs
leveled binary tree = directed tree with a binary search tree labeling and an increasing labeling



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees

The sylvester correspondence = permutations \longmapsto leveled binary trees.


Exm: permutation 6275134



Permutations to leveled binary trees


The sylvester correspondence = permutations \longmapsto leveled binary trees.

Exm: permutation 6275134




Cambrian trees

Cambrian tree = directed and labeled tree such that




increasing tree = directed and labeled tree such that labels increase along arcs
leveled Cambrian tree = directed tree with a Cambrian labeling and an increasing labeling



Cambrian trees and triangulations

Cambrian trees are dual to triangulations of polygons




signature \longleftrightarrow vertices above or below $[0, 8]$
node $j \longleftrightarrow$ triangle $i < j < k$

For any signature ε , there are $C_n = \frac{1}{n+1} \binom{2n}{n}$ ε -Cambrian trees

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longleftrightarrow leveled Cambrian tree.

Exm: signed permutation $\underline{2}\bar{7}5\underline{1}\bar{3}\bar{4}\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longmapsto leveled Cambrian tree.

Exm: signed permutation $2\bar{7}51\bar{3}4\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longmapsto leveled Cambrian tree.

Exm: signed permutation $2\bar{7}51\bar{3}4\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longmapsto leveled Cambrian tree.

Exm: signed permutation $2\bar{7}51\bar{3}4\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longleftrightarrow leveled Cambrian tree.

Exm: signed permutation $2\bar{7}51\bar{3}4\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longleftrightarrow leveled Cambrian tree.

Exm: signed permutation $2\bar{7}51\bar{3}4\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longleftrightarrow leveled Cambrian tree.

Exm: signed permutation $2\bar{7}51\bar{3}4\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longleftrightarrow leveled Cambrian tree.

Exm: signed permutation $\underline{2}\bar{7}5\underline{1}\bar{3}\bar{4}\bar{6}$




Reading. Cambrian lattices. 2006
Lange-Pilaud. Associahedra via spines. 2015

Signed permutations to Cambrian trees

Cambrian correspondence = signed permutation \longleftrightarrow leveled Cambrian tree.

Exm: signed permutation $\underline{2}\bar{7}5\underline{1}\bar{3}\bar{4}\bar{6}$



$\mathbf{P}(\tau)$ = P-symbol of τ = Cambrian tree produced by the Cambrian corresp.
 $\mathbf{Q}(\tau)$ = Q-symbol of τ = increasing tree produced by the Cambrian corresp.

(analogous to the Robinson-Schensted algorithm)

The Cambrian congruence

ε -Cambrian congruence = transitive closure of the rewriting rules


$\cdots ac \cdots \underline{b} \cdots \equiv_{\varepsilon} \cdots ca \cdots \underline{b} \cdots$ if $a < b < c$ and $\varepsilon_b = -$

$\cdots \bar{b} \cdots ac \cdots \equiv_{\varepsilon} \cdots \bar{b} \cdots ca \cdots$ if $a < b < c$ and $\varepsilon_b = +$


where a, b, c are elements of $[n]$.

Proposition [reformulating Reading 2006]

$$\tau \equiv_{\varepsilon} \tau' \iff \mathbf{P}(\tau) = \mathbf{P}(\tau')$$



$\underline{2}\bar{7}51\bar{3}4\bar{6}$




$\bar{7}2513\bar{4}6$

Cambrian trees to signed permutations


Proposition [reformulating Reading 2006]

$$\mathbf{P}^{-1}(T) = \mathcal{L}(T)$$




$$\mathbf{P}^{-1}(T) = \mathcal{L}(T) = \{\underline{2}1\underline{3}\overline{7}54\overline{6}, \underline{2}1\overline{7}354\overline{6}, \underline{2}1\overline{7}5\underline{3}4\overline{6}, \underline{2}\overline{7}1\overline{3}54\overline{6}, \underline{2}\overline{7}15\overline{3}4\overline{6}, \\ \underline{2}\overline{7}5134\overline{6}, \underline{7}21\overline{3}54\overline{6}, \underline{7}21534\overline{6}, \overline{7}25134\overline{6}, \overline{7}52134\overline{6}\}$$

Rotations and flips

Rotation on Cambrian trees \longleftrightarrow flips on triangulations.

Rotation and Cambrian lattices

Rotation operation preserves Cambrian trees:





increasing rotation = rotation of edge $i \rightarrow j$ where $i < j$

Proposition [reformulating Reading 2006]

The transitive closure of the increasing rotation graph is the Cambrian lattice.
 \mathbf{P} defines a lattice homomorphism from the weak order to the Cambrian lattice.

Rotations and Cambrian lattices



1 Combinatorics

- Binary trees
- Cambrian trees
- Cambrian lattices

2 Algebra

- FQSym
- The Cambrian algebra


Two products on permutations

For $\tau \in \mathfrak{S}_n$ and $\tau' \in \mathfrak{S}_{n'}$ with $a \in [n], b \in [n']$, define


- shifted concatenation $\tau\bar{\tau}' = [\tau(1), \dots, \tau(n), \tau'(1) + n, \dots, \tau'(n') + n] \in \mathfrak{S}_{n+n'}$
- shifted shuffle product $\tau \sqcup \tau' = au \sqcup bv = a(u \sqcup bv) + (b + |au|)(au \sqcup v)$
- convolution product $\tau * \tau' = (\tau^{-1} \sqcup \tau'^{-1})^{-1}$

When we compute products of permutations, there is no multiplicities so we can consider that the output of the shuffle is a set of permutations.


$$\begin{aligned} 12 \sqcup 231 &= \{12453, 14253, 14523, 14532, 41253, 41523, 41532, 45123, 45132, 45312\} \\ 12 * 231 &= \{12453, 13452, 14352, 15342, 23451, 24351, 25341, 34251, 35241, 45231\} \end{aligned}$$



concatenation



shuffle



convolution

The Malvenuto-Reutenauer algebra

The Malvenuto-Reutenauer algebra = Hopf algebra FQSym with basis $(\mathbb{F}_\tau)_{\tau \in \mathfrak{S}}$ and where

$$\mathbb{F}_\tau \cdot \mathbb{F}_{\tau'} = \sum_{\sigma \in \tau \sqcup \tau'} \mathbb{F}_\sigma \quad \text{and} \quad \Delta \mathbb{F}_\sigma = \sum_{\sigma \in \tau \star \tau'} \mathbb{F}_\tau \otimes \mathbb{F}_{\tau'}$$

Malvenuto-Reutenauer. Duality between Quasi-Symmetric functions and the Solomon Descent Algebra. 1995

The Malvenuto-Reutenauer algebra

The Malvenuto-Reutenauer algebra = Hopf algebra FQSym with basis $(\mathbb{F}_\tau)_{\tau \in \mathfrak{S}}$ and where

$$\mathbb{F}_\tau \cdot \mathbb{F}_{\tau'} = \sum_{\sigma \in \tau \sqcup \tau'} \mathbb{F}_\sigma \quad \text{and} \quad \Delta \mathbb{F}_\sigma = \sum_{\sigma \in \tau \star \tau'} \mathbb{F}_\tau \otimes \mathbb{F}_{\tau'}$$

Malvenuto-Reutenauer. Duality between Quasi-Symmetric functions and the Solomon Descent Algebra. 1995

Definition: Combinatorial Hopf Algebras

A Combinatorial Hopf Algebra = combinatorial vector space \mathcal{B} endowed with

$$\begin{aligned} \text{product } \cdot &: \mathcal{B} \otimes \mathcal{B} \rightarrow \mathcal{B} \\ \text{coproduct } \Delta &: \mathcal{B} \rightarrow \mathcal{B} \otimes \mathcal{B} \end{aligned}$$


which are “compatible”, i.e.,

$$\Delta(f \cdot g) = \Delta(f) \cdot \Delta(g)$$


Two products on signed permutations

For signed permutations:


$$\begin{aligned}\bar{1}2 \sqcup \underline{2}3\bar{1} &= \{\bar{1}245\bar{3}, \bar{1}425\bar{3}, \bar{1}45\bar{2}\bar{3}, \bar{1}45\bar{3}2, \bar{4}125\bar{3}, 4\bar{1}5\bar{2}\bar{3}, 4\bar{1}5\bar{3}2, 45\bar{1}\bar{2}\bar{3}, 45\bar{1}\bar{3}2, 45\bar{3}\bar{1}2\}, \\ \bar{1}2 \star \underline{2}3\bar{1} &= \{\bar{1}245\bar{3}, \bar{1}3452, \bar{1}4352, \bar{1}5342, \bar{2}345\bar{1}, \bar{2}435\bar{1}, \bar{2}534\bar{1}, \bar{3}425\bar{1}, \bar{3}524\bar{1}, \bar{4}523\bar{1}\}.\end{aligned}$$



concatenation



shuffle



convolution

Signed analog of Malvenuto-Reutenauer [Novelli, Thibon 2010]

FQSym_{\pm} = Hopf algebra with basis $(\mathbb{F}_\tau)_{\tau \in \mathfrak{S}_{\pm}}$ and where

$$\mathbb{F}_\tau \cdot \mathbb{F}_{\tau'} = \sum_{\sigma \in \tau \sqcup \tau'} \mathbb{F}_\sigma \quad \text{and} \quad \Delta \mathbb{F}_\sigma = \sum_{\sigma \in \tau \star \tau'} \mathbb{F}_\tau \otimes \mathbb{F}_{\tau'}$$

The Cambrian algebra as a subalgebra of FQSym_\pm

The Cambrian algebra = subspace Camb of FQSym_\pm generated by

$$\mathbb{P}_T := \sum_{\substack{\tau \in \mathfrak{S}_\pm \\ P(\tau) = T}} \mathbb{F}_\tau = \sum_{\tau \in \mathcal{L}(T)} \mathbb{F}_\tau,$$

for all Cambrian trees T .



$$\begin{aligned} \mathbb{P} = & \mathbb{F}_{\underline{21}\bar{3}\bar{7}\underline{54}\bar{6}} + \mathbb{F}_{\underline{21}\bar{7}\bar{3}\underline{54}\bar{6}} + \mathbb{F}_{\underline{21}\bar{7}\bar{5}\bar{3}\underline{46}} + \mathbb{F}_{\bar{2}\bar{7}\bar{1}\bar{3}\underline{54}\bar{6}} + \mathbb{F}_{\bar{2}\bar{7}\bar{1}\bar{5}\bar{3}\underline{46}} \\ & + \mathbb{F}_{\bar{2}\bar{7}\bar{5}\bar{1}\bar{3}\underline{46}} + \mathbb{F}_{\bar{7}\underline{21}\bar{3}\underline{54}\bar{6}} + \mathbb{F}_{\bar{7}\bar{2}\bar{1}\bar{5}\bar{3}\underline{46}} + \mathbb{F}_{\bar{7}\bar{2}\bar{5}\bar{1}\bar{3}\underline{46}} + \mathbb{F}_{\bar{7}\bar{5}\bar{2}\bar{1}\bar{3}\underline{46}} \end{aligned}$$

Theorem [C.-Pilaud]

Camb is a Hopf subalgebra of FQSym_\pm .

(i.e., the Cambrian congruence is “compatible” with the product and coproduct in FQSym_\pm)

GAME: Explain the product and coproduct directly on the Cambrian trees...

Product in the Cambrian algebra

$$\begin{aligned}
 \mathbb{P} \text{ } \begin{array}{c} \diagup \\ \diagdown \end{array} \cdot \mathbb{P} \text{ } \begin{array}{c} \diagup \\ \diagdown \end{array} &= \mathbb{F}_{\underline{1}\bar{2}} \cdot (\mathbb{F}_{\bar{2}\underline{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\underline{1}}) \\
 &= \left(\begin{array}{l} \mathbb{F}_{\underline{1}\bar{2}\bar{4}\underline{3}\bar{5}} + \mathbb{F}_{\underline{1}\bar{2}\bar{4}\bar{5}\underline{3}} + \mathbb{F}_{\bar{1}\bar{4}\bar{2}\underline{3}\bar{5}} \\ + \mathbb{F}_{\underline{1}\bar{4}\bar{2}\underline{5}\bar{3}} + \mathbb{F}_{\bar{1}\bar{4}\bar{5}\bar{2}\underline{3}} + \mathbb{F}_{\bar{4}\bar{1}\bar{2}\bar{3}\bar{5}} \\ + \mathbb{F}_{\bar{4}\bar{1}\bar{2}\bar{5}\bar{3}} + \mathbb{F}_{\bar{4}\bar{1}\bar{5}\bar{2}\bar{3}} + \mathbb{F}_{\bar{4}\bar{5}\bar{1}\bar{2}\bar{3}} \end{array} \right) + \left(\begin{array}{l} \mathbb{F}_{\bar{1}\bar{4}\bar{3}\bar{2}\bar{5}} + \mathbb{F}_{\bar{1}\bar{4}\bar{3}\bar{5}\bar{2}} \\ + \mathbb{F}_{\bar{1}\bar{4}\bar{5}\bar{3}\bar{2}} + \mathbb{F}_{\bar{4}\bar{1}\bar{3}\bar{2}\bar{5}} \\ + \mathbb{F}_{\bar{4}\bar{1}\bar{3}\bar{5}\bar{2}} + \mathbb{F}_{\bar{4}\bar{1}\bar{5}\bar{3}\bar{2}} \\ + \mathbb{F}_{\bar{4}\bar{5}\bar{1}\bar{3}\bar{2}} \end{array} \right) + \left(\begin{array}{l} \mathbb{F}_{\bar{4}\bar{3}\bar{1}\bar{2}\bar{5}} + \mathbb{F}_{\bar{4}\bar{3}\bar{1}\bar{5}\bar{2}} \\ + \mathbb{F}_{\bar{4}\bar{3}\bar{5}\bar{1}\bar{2}} + \mathbb{F}_{\bar{4}\bar{5}\bar{3}\bar{1}\bar{2}} \end{array} \right) \\
 &= \mathbb{P} \text{ } \begin{array}{c} \diagup \\ \diagdown \end{array} \quad + \quad \mathbb{P} \text{ } \begin{array}{c} \diagup \\ \diagdown \end{array} \quad + \quad \mathbb{P} \text{ } \begin{array}{c} \diagup \\ \diagdown \end{array}
 \end{aligned}$$

Proposition [C.-Pilaud]

For any Cambrian trees T and T' ,

$$\mathbb{P}_T \cdot \mathbb{P}_{T'} = \sum_{\substack{T' \nearrow \overline{T'} \\ T \searrow \overline{T}}} \mathbb{P}_S$$

Product in the Cambrian algebra

Proposition [C.-Pilaud]

For any Cambrian trees T and T' ,

$$\mathbb{P}_T \cdot \mathbb{P}_{T'} = \sum_{\substack{T \nearrow \overline{T'} \\ S \leq_{\text{Camb}} T' \nwarrow \overline{T}}} \mathbb{P}_S$$

- For every Cambrian tree T , $\mathcal{L}(T)$ is an interval of the weak order.

Product in the Cambrian algebra

Proposition [C.-Pilaud]

For any Cambrian trees T and T' ,

$$\mathbb{P}_T \cdot \mathbb{P}_{T'} = \sum_{\substack{T \nearrow T' \\ S \leq_{\text{Camb}} T' \nwarrow T}} \mathbb{P}_S$$

- For every Cambrian tree T , $\mathcal{L}(T)$ is an interval of the weak order.
- $[\sigma, \tau] \boxplus [\sigma', \tau'] = [\sigma\overline{\sigma'}, \overline{\tau'}\tau]$.

Product in the Cambrian algebra

Proposition [C.-Pilaud]

For any Cambrian trees T and T' ,

$$\mathbb{P}_T \cdot \mathbb{P}_{T'} = \sum_{\substack{T' \\ \nearrow \overline{T'} \\ T}} \leq_{\text{Camb}} S \leq_{\text{Camb}} \sum_{\substack{T \\ \searrow \overline{T'} \\ \overline{T'}}} \mathbb{P}_S$$

- For every Cambrian tree T , $\mathcal{L}(T)$ is an interval of the weak order.
- $[\sigma, \tau] \boxplus [\sigma', \tau'] = [\sigma\overline{\sigma'}, \overline{\tau'}\tau]$.
- $\mathcal{L}(T) = [\sigma, \tau], \mathcal{L}(T') = [\sigma', \tau']$

$$\mathbb{P}(\sigma\overline{\sigma'}) = \begin{array}{c} \nearrow \overline{T'} \\ T \end{array} \quad \mathbb{P}(\overline{\tau'}\tau) = \begin{array}{c} T \\ \searrow \overline{T'} \end{array}$$

Product in the Cambrian algebra

Proposition [C.-Pilaud]

For any Cambrian trees T and T' ,

$$\mathbb{P}_T \cdot \mathbb{P}_{T'} = \sum_{\substack{T' \\ \nearrow \overline{T'} \\ T}} \leq_{\text{Camb}} S \leq_{\text{Camb}} \sum_{\substack{T \\ \searrow \overline{T'} \\ \overline{T'}}} \mathbb{P}_S$$

- For every Cambrian tree T , $\mathcal{L}(T)$ is an interval of the weak order.
- $[\sigma, \tau] \boxplus [\sigma', \tau'] = [\sigma\overline{\sigma'}, \overline{\tau'}\tau]$.
- $\mathcal{L}(T) = [\sigma, \tau], \mathcal{L}(T') = [\sigma', \tau']$

$$\mathbb{P}(\sigma\overline{\sigma'}) = \begin{array}{c} \nearrow \overline{T'} \\ T \end{array} \quad \mathbb{P}(\overline{\tau'}\tau) = \begin{array}{c} T \\ \searrow \overline{T'} \end{array}$$

- Permutations in $[\sigma\overline{\sigma'}, \overline{\tau'}\tau]$ form a union of Cambrian classes because Camb is a subalgebra of FQSym $_{\pm}$.

Product in the Cambrian algebra

Proposition [C.-Pilaud]

For any Cambrian trees T and T' ,

$$\mathbb{P}_T \cdot \mathbb{P}_{T'} = \sum_{\substack{T' \\ \nearrow \overline{T'} \\ T}} \leq_{\text{Camb}} S \leq_{\text{Camb}} \sum_{\substack{T \\ \searrow \overline{T'} \\ \overline{T'}}} \mathbb{P}_S$$

- For every Cambrian tree T , $\mathcal{L}(T)$ is an interval of the weak order.
- $[\sigma, \tau] \sqcup [\sigma', \tau'] = [\sigma\overline{\sigma'}, \overline{\tau'}\tau]$.
- $\mathcal{L}(T) = [\sigma, \tau], \mathcal{L}(T') = [\sigma', \tau']$

$$\mathbf{P}(\sigma\overline{\sigma'}) = \begin{array}{c} \nearrow \overline{T'} \\ T \end{array} \quad \mathbf{P}(\overline{\tau'}\tau) = \begin{array}{c} T \\ \searrow \overline{T'} \end{array}$$

- Permutations in $[\sigma\overline{\sigma'}, \overline{\tau'}\tau]$ form a union of Cambrian classes because Camb is a subalgebra of FQSym_{\pm} .
- These trees form an interval because \mathbf{P} is a lattice homomorphism.

Coproduct in the Cambrian algebra


$$\begin{aligned}
 \Delta \mathbb{P} &= \Delta(\mathbb{F}_{\bar{2}\bar{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\bar{1}}) \\
 &= 1 \otimes (\mathbb{F}_{\bar{2}\bar{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\bar{1}}) + \mathbb{F}_{\bar{1}} \otimes \mathbb{F}_{\bar{1}\bar{2}} + \mathbb{F}_{\bar{1}} \otimes \mathbb{F}_{\bar{2}\bar{1}} + \mathbb{F}_{\bar{2}\bar{1}} \otimes \mathbb{F}_{\bar{1}} + \mathbb{F}_{\bar{1}\bar{2}} \otimes \mathbb{F}_{\bar{1}} + (\mathbb{F}_{\bar{2}\bar{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\bar{1}}) \otimes 1 \\
 &= 1 \otimes \mathbb{P} + \mathbb{P} \otimes \mathbb{P} \\
 &= 1 \otimes \mathbb{P} + \mathbb{P} \otimes (\mathbb{P} \cdot \mathbb{P}) + \mathbb{P} \otimes \mathbb{P} + \mathbb{P} \otimes 1.
 \end{aligned}$$

Proposition [C.-Pilaud]

For any Cambrian tree S ,

$$\Delta \mathbb{P}_S = \sum_{\gamma} \left(\prod_{T \in B(S, \gamma)} \mathbb{P}_T \right) \otimes \left(\prod_{T' \in A(S, \gamma)} \mathbb{P}_{T'} \right)$$

where γ runs over all cuts of S , and $A(S, \gamma)$ and $B(S, \gamma)$ denote the Cambrian forests above and below γ respectively



Coproduct in the Cambrian algebra


$$\begin{aligned}
 \Delta \mathbb{P} \text{ (Diagram)} &= \Delta (\mathbb{F}_{\bar{2}\bar{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\bar{1}}) \\
 &= 1 \otimes (\mathbb{F}_{\bar{2}\bar{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\bar{1}}) + \mathbb{F}_{\bar{1}} \otimes \mathbb{F}_{\bar{1}\bar{2}} + \mathbb{F}_{\bar{1}} \otimes \mathbb{F}_{\bar{2}\bar{1}} + \mathbb{F}_{\bar{2}\bar{1}} \otimes \mathbb{F}_{\bar{1}} + \mathbb{F}_{\bar{1}\bar{2}} \otimes \mathbb{F}_{\bar{1}} + (\mathbb{F}_{\bar{2}\bar{1}\bar{3}} + \mathbb{F}_{\bar{2}\bar{3}\bar{1}}) \otimes 1 \\
 &= 1 \otimes \mathbb{P} \text{ (Diagram)} + \mathbb{P} \otimes \mathbb{P} \text{ (Diagram)} \otimes 1 \\
 &= 1 \otimes \mathbb{P} \text{ (Diagram)} + \mathbb{P} \otimes (\mathbb{P} \cdot \mathbb{P}) \text{ (Diagram)} + \mathbb{P} \otimes \mathbb{P} \text{ (Diagram)} \otimes 1.
 \end{aligned}$$

Proposition [C.-Pilaud]

For any Cambrian tree S ,

$$\Delta \mathbb{P}_S = \sum_{\gamma} \left(\prod_{T \in B(S, \gamma)} \mathbb{P}_T \right) \otimes \left(\prod_{T' \in A(S, \gamma)} \mathbb{P}_{T'} \right)$$

where γ runs over all cuts of S , and $A(S, \gamma)$ and $B(S, \gamma)$ denote the Cambrian forests above and below γ respectively



Extensions

- Study of multiplicative basis.
- Hopf algebra on twin Cambrian trees.
- Hopf algebra on Schröder-Cambrian trees.

